

Report IRB-StP-GR-240696

# The Simplest Exact Solutions in the LTB Model

*by*

**Alexander Gromov**

*St. Petersburg State Technical University*

*Faculty of Technical Cybernetics, Dept. of Computer Science*

*29, Polytechnicheskaya str. St.-Petersburg, 195251, Russia*

and

*Istituto per la Ricerca di Base*

*Castello Principe Pignatelli del Comune di Monteroduni*

*I-86075 Monteroduni(IS), Molise, Italia*

*e-mail: gromov@natus.stud.pu.ru*

## Abstract

The rules of calculating three undetermined functions which defined a solution in the LTB model are used to study the class of exact nonhomogeneous models with  $f^2(\mu) = 1$ ,  $\Lambda = 0$ . The parameter  $\nu(\mu)$  defined the difference between LTB and FRW models is found out and the limit transformation to the FRW model is shown. The initial conditions are present through density and Hubble function at the moment of time  $\tau = 0$ . Two criteria of homogeneous of matter distribution are studied. The asymptotic of the present solution for  $\tau \rightarrow +\infty$  is studied.

PACS number(s):98.80

# 1 The Introduction

This article is dedicated to studying of the simplest solution in the LTB model and based on the results are obtained in [1],[2] and [3] and presented shortly in this section. The LTB model is studied here as the Cauchy problem for the PDE [1]:

$$e^{3\omega(\mu,\tau)/2} \left( \frac{\dot{\omega}^2(\mu,\tau)}{2} - \frac{2}{3}\Lambda \right) + 2e^{\omega(\mu,\tau)/2} [1 - f^2(\mu)] = F(\mu) \quad (1)$$

with initial conditions

$$\left. \begin{aligned} \omega(\mu, \tau)|_{\tau=0} &= \omega_0(\mu), & \dot{\omega}(\mu, \tau)|_{\tau=0} &= \dot{\omega}_0(\mu), \\ \ddot{\omega}(\mu, \tau)|_{\tau=0} &= \ddot{\omega}_0(\mu) \end{aligned} \right\} \quad (2)$$

and constants

$$\left. \begin{aligned} \omega(\mu, 0)|_{\mu=0} &= \omega_0(0), & \dot{\omega}(\mu, 0)|_{\mu=0} &= \dot{\omega}_0(0), \\ \ddot{\omega}(\mu, 0)|_{\mu=0} &= \ddot{\omega}_0(0), & \Lambda, \end{aligned} \right\} \quad (3)$$

where  $\mu$  and  $\tau$  are dimensionless co-moving coordinates, corresponding to Lagrangian coordinate and time,  $\omega(\mu, \tau)$  is the metrical function,  $\dot{\omega} = \frac{\partial \omega}{\partial \tau}$ ,  $\Lambda$  is the cosmological constant,

$$f^2(\mu) - 1 = e^{\omega_0(\mu)} \left( \ddot{\omega}_0(\mu) + \frac{3}{4}\dot{\omega}_0^2(\mu) - \Lambda \right), \quad (4)$$

$$F(\mu) = e^{3\omega_0(\mu)/2} \left( \frac{\dot{\omega}_0^2(\mu)}{2} - \frac{2}{3}\Lambda \right) + 2e^{\omega_0(\mu)/2} [1 - f^2(\mu)]. \quad (5)$$

The law of the density has the form

$$\begin{aligned} 8\pi\delta(\mu, \tau) &= \frac{e^{\frac{3}{2}[\omega_0(\mu) - \omega(\mu, \tau)]}}{\omega'(\mu, \tau)} \times \\ &\left\{ 3[\omega_0(\mu)]' \left[ -\ddot{\omega}_0(\mu) - \frac{1}{2}\dot{\omega}_0^2(\mu) + \frac{\Lambda}{6} \right] - 2[\ddot{\omega}_0(\mu)]' - 2\dot{\omega}_0(\mu) [\dot{\omega}_0(\mu)]' \right\} \end{aligned} \quad (6)$$

The Habb function is defined by the expression

$$h(\mu, \tau) = \frac{\dot{R}'(\mu, \tau)}{R'(\mu, \tau)} = \frac{\partial \ln R'(\mu, \tau)}{\partial \tau}. \quad (7)$$

This article studies the simplest solution in the LTB model defining by the initial conditions

$$f^2(\mu) = 1, \quad \Lambda = 0. \quad (8)$$

For the first time this solution has obtained by Bonnor [2]. The Bonnor's result is wrote for the Bonnor's function

$$R(\mu, \tau) = e^{\omega(\mu, \tau)/2},$$

which has the sense of Euler coordinate [4], [5]:

$$R(\mu, \tau) = \frac{9^{1/3}}{2} \left[ \tau \sqrt{\Psi(\mu)} + \beta(\mu) \right]^{2/3}, \quad (9)$$

where  $\Psi(\mu)$  and  $\beta(\mu)$  are arbitrary functions.

This article is devoted to study the same problem through the metrical function and initial conditions for it.

## 2 The $f^2(\mu) = 1$ , $\Lambda = 0$ Solution

The initial conditions (8) generate the simplest analytical solution. The condition  $f^2(\mu) = 1$ , as it goes from (4), correlates the initial conditions as follows:

$$\ddot{\omega}_0(\mu) + \frac{3}{4}\dot{\omega}_0^2(\mu) = 0. \quad (10)$$

Due to this reason, the general number of the initial conditions (2) is decreased by one unit.

We start with the solution of the equation (1). It follows from (1) that:

$$\left(\frac{\partial e^{\omega(\mu,\tau)/2}}{\partial \tau}\right)^2 = f^2(\mu) - 1 + \frac{1}{2}F(\mu)e^{-\omega(\mu,\tau)/2} + \frac{\Lambda}{3}e^{\omega}. \quad (11)$$

In the general case the function  $F(\mu)$  is defined by the equation (5). With help of (8) and (10) it takes the form

$$F(\mu) = \frac{\dot{\omega}_0^2(\mu)}{2}e^{3\omega_0(\mu)/2}$$

and the equation (11) with conditions (8) becomes as follows:

$$e^{3\omega_0(\mu)/4} \frac{\partial e^{3\omega(\mu,\tau)/4}}{\partial \tau} = \pm \frac{3}{4}\dot{\omega}_0(\mu). \quad (12)$$

We obtain by the integration of (12)

$$e^{\frac{3}{4}[\omega(\mu,\tau) - \omega_0(\mu)]} = \pm \frac{3}{4}\dot{\omega}_0(\mu)\tau + \mathbf{F}(\mu).$$

The function  $\mathbf{F}(\mu)$  is found out from the initial conditions then time  $\tau = 0$ :

$$\mathbf{F}(\mu) = 1.$$

The solution now is:

$$e^{\frac{3}{4}[\omega(\mu,\tau) - \omega_0(\mu)]} = 1 \pm \frac{3}{4}\dot{\omega}_0(\mu)\tau. \quad (13)$$

In the Bonnor's notation the equation (13) is readied:

$$\frac{R(\mu,\tau)}{R_0(\mu)} = \left(1 \pm \frac{3}{4}\dot{\omega}_0(\mu)\tau\right)^{2/3}, \quad (14)$$

where the upper sign correspond to the expansion from the initial condition into the infinity and lower sign correspond to the collapse from one.

The collapse solutions contains the characteristic time of collapse depending on the initial mass coordinate: if the particle has the coordinate  $\mu$  at the moment  $\tau = 0$ , the time of collapse is equal to

$$\bar{\tau}(\mu) = \frac{4}{3\dot{\omega}_0(\mu)}. \quad (15)$$

From this moment and up to the end of this section the dependence of  $\omega$  from  $\mu$  and  $\tau$  will be omitted. We obtain the density corresponding to this solution by the substitution (13) into (6). Let's denote

$$\nu = \frac{[\dot{\omega}_0]'}{[\omega_0]'}. \quad (16)$$

So, the density is:

$$8\pi\delta(\mu,\tau) = \frac{\dot{\omega}_0}{\left(1 \pm \frac{3}{4}\dot{\omega}_0\tau\right)^2} \cdot \frac{\nu + \frac{3}{4}\dot{\omega}_0}{1 \pm \frac{\frac{3}{4}\dot{\omega}_0\tau}{\nu}}, \quad (17)$$

Using the definition (7) we find out the Hubble's function for the solution (13):

$$h(\mu, \tau) = \pm \frac{\frac{3}{4}\dot{\omega}_0 + \nu}{1 \pm \left(\frac{3}{4}\dot{\omega}_0 + \nu\right)\tau} \mp \frac{1}{4} \cdot \frac{\dot{\omega}_0}{1 \pm \frac{3}{4}\dot{\omega}_0\tau} \quad (18)$$

Knowing the law of the density and Hubble's function we obtain now the formulism for the cosmological parameter  $\Omega$ :

$$\Omega = \frac{\rho}{\rho_c}, \quad \text{where} \quad \rho_c = \frac{3H_0^2}{8\pi G},$$

where  $H_0$  and  $\rho$  mean the dimension Hubble function and the density at the moment of the observation. Let's assume that this moment is  $\tau = 0$ . The density, the critical density and Hubble function at this moment are:

$$8\pi\delta(\mu, 0) = \dot{\omega}_0 \left( \frac{3}{4}\dot{\omega}_0 + \nu \right), \quad (19)$$

$$\delta_c(\mu, 0) = \frac{3\tilde{H}^2}{8\pi G\rho(0, 0)} h^2(\mu, 0),$$

$$h(\mu, 0) = \pm \left( \frac{1}{2}\dot{\omega}_0 + \nu \right), \quad (20)$$

$$\Omega(\mu, 0) = \frac{G\rho_0}{3\tilde{H}^2} \dot{\omega}_0 \frac{\frac{3}{4}\dot{\omega}_0 + \nu}{\left(\frac{\dot{\omega}_0}{2} + \nu\right)^2}.$$

These functions dependent on time as follow:

$$\delta_c(\mu, \tau) = \frac{3\tilde{H}^2}{8\pi G\rho_0} \left[ \pm \frac{\frac{3}{4}\dot{\omega}_0 + \nu}{1 \pm \left(\frac{3}{4}\dot{\omega}_0 + \nu\right)\tau} \mp \frac{1}{4} \cdot \frac{\dot{\omega}_0}{1 \pm \frac{3}{4}\dot{\omega}_0\tau} \right]^2,$$

$$\Omega(\mu, \tau) = \frac{16G\rho_0}{3\tilde{H}^2} \dot{\omega}_0 \left( \frac{3}{4}\dot{\omega}_0 + \nu \right) \times \frac{(1 \pm \frac{3}{4}\dot{\omega}_0\tau) [1 \pm (\frac{3}{4}\dot{\omega}_0 + \nu)\tau]}{\{\pm 4(1 \pm \frac{3}{4}\dot{\omega}_0\tau) (\frac{3}{4}\dot{\omega}_0 + \nu) \mp \dot{\omega}_0 [1 \pm (\frac{3}{4}\dot{\omega}_0 + \nu)\tau]\}^2}.$$

### 3 The FRW Model

Every nonhomogeneous solution of the LTB model should definitely include the FRW model as a limited case when the density and Hubble function are not dependent on space coordinate for all moments of time. Let's study in which case the present solution is reduced to the FRW model? Only one condition satisfies this request:

$$\omega_0(\mu) = \text{const} \quad \text{so,} \quad \nu(\mu) = 0. \quad (21)$$

It goes from (17), (18) and (21) that in this case

$$8\pi\delta(\tau) = \frac{3}{4} \left( \frac{\dot{\omega}_0}{1 \pm \frac{3}{4}\dot{\omega}_0\tau} \right)^2, \quad (22)$$

$$h(\tau) = \pm \frac{1}{2} \cdot \frac{\dot{\omega}_0}{1 \pm \frac{3}{4}\dot{\omega}_0\tau}.$$

From these equations it goes:

$$8\pi\delta(\tau) = 3h^2(\tau). \quad (23)$$

We obtain the meaning of the  $\omega_0$  by substitution  $\tau = 0$  in (22):

$$\dot{\omega}_0 = \pm 2\sqrt{\frac{8\pi}{3}}. \quad (24)$$

For the collapse solution, in accordance with (15) all matter fall down to the centre simultaneously at time

$$\bar{\tau}(\mu) = \frac{1}{\sqrt{6\pi}}. \quad (25)$$

$\nu(\mu) = 0$  together with (24) give

$$\Omega = \frac{G\rho_0}{\tilde{H}^2}.$$

### 4 The Initial Conditions and Criteria of Homogeneous

We are able now to construct here the initial conditions for the  $f^2 = 1$ ,  $\Lambda = 0$  solution. To do this, we assume the functions  $\delta(\mu, 0)$  and  $h(\mu, 0)$  are given and express the initial conditions  $\omega_0(\mu)$  and  $\dot{\omega}_0(\mu)$  through  $\delta(\mu, 0)$  and  $h(\mu, 0)$ . From (19) and (20) we find out:

$$\dot{\omega}_0(\mu) = 2\epsilon h(\mu, 0) \pm \sqrt{4h^2(\mu, 0) + 32\pi\delta(\mu, 0)} \quad (26)$$

and

$$\nu(\mu) = \frac{8\pi\delta(\mu, 0)}{2\epsilon h(\mu, 0) \pm \sqrt{4h^2(\mu, 0) + 32\pi\delta(\mu, 0)}} - \frac{3}{4}\{2\epsilon h(\mu, 0) \pm \sqrt{4h^2(\mu, 0) + 32\pi\delta(\mu, 0)}\} \quad (27)$$

where  $\epsilon = -1$  for the expansion solution and  $\epsilon = +1$  for the collapse one. It was shown in the section (3) that the general solution is reduced to the FRW by the equality  $\nu(\mu) = 0$ . Solving the equation  $\nu(\mu) = 0$  with  $\nu(\mu)$  from (27) we obtain for  $\frac{8\pi\delta}{h^2}$  the quadratic equation which has two solutions:

$$\frac{8\pi\delta}{h^2} = 0 \quad \frac{8\pi\delta}{h^2} = 3.$$

First of them is the solution for the empty space and the second one is the solution for the FRW model.

We will study now two criteria of homogeneity of the matter distribution. First of them is the difference

$$8\pi\delta(\mu, 0) - 3h^2(\mu, 0) = -\nu(\mu) [2\dot{\omega}_0(\mu) + 3\nu(\mu)] \quad (28)$$

The right part of this equation is equal to zero in two cases:

$$\nu(\mu) = \frac{[\dot{\omega}_0(\mu)]'}{[\omega_0(\mu)]'} = 0 \quad (29)$$

and

$$\nu(\mu) = \frac{[\dot{\omega}_0(\mu)]'}{[\omega_0(\mu)]'} = -\frac{2}{3}\dot{\omega}_0(\mu). \quad (30)$$

Solving the equation (30) as differential equation for the function  $\dot{\omega}_0(\mu)$  (initial conditions) we find out the correlation  $\dot{\omega}_0(\mu)$  and  $\omega_0(\mu)$ :

$$\dot{\omega}_0(\mu) = \text{const} \cdot e^{2\omega_0(\mu)/3}$$

But the solution (30) do not assume the density  $\delta(\mu, 0) = \text{const}$  excluding the case (29). This example shows the existence of such initial conditions that the equality (28) is satisfied for density not equal to constant at the moment of time  $\tau = 0$ .

The second criterium is the ratio  $\frac{\delta_{LTB}}{\delta_{FRW}}$  where  $\delta_{LTB}$  is from (17) and  $\delta_{FRW}$  is from (22):

$$\frac{\delta_{LTB}}{\delta_{FRW}} = \frac{4}{3\dot{\omega}_0} \left( 1 \pm \frac{3}{4}\dot{\omega}_0\tau \right) \frac{\frac{3}{4}\dot{\omega}_0 + \nu}{1 \pm \left( \frac{3}{4}\dot{\omega}_0 + \nu \right) \tau} \quad (31)$$

In accordance with the FRW solution the equality  $\nu(\mu) = 0$  implies the equality  $\frac{\delta_{LTB}}{\delta_{FRW}} = 1$ .

## 5 The Property of the Solution for

$$\tau \rightarrow +\infty$$

Let's study now the expansion nonhomogeneous model with presented in the section (2) for time

$$\tau \gg \bar{\tau}(\mu) = \frac{4}{3\dot{\omega}_0(\mu)}. \quad (32)$$

The sense of time  $\bar{\tau}$  is shown by (15) for collapse solution. In the expansion model it is characteristic value connected with initial conditions:  $\mu$  is the initial position of the particle at the moment of time  $\tau = 0$ . In the limit (32) the density and Hubble's function are not dependent on  $\mu$ -coordinate but only on time:

$$8\pi\delta(\tau) \rightarrow \frac{4}{3\tau^2},$$

$$h(\tau) \rightarrow \frac{2}{3\tau},$$

so, the follows equation takes a place in the limit (32):

$$8\pi \cdot \lim_{\tau \gg \bar{\tau}(\mu)} \delta(\tau) = 3 \cdot \lim_{\tau \gg \bar{\tau}(\mu)} h^2(\tau)$$

This property allows to name the solution (13) - (14) of the nonhomogeneous model in the area (32) by FRW-limit of LTB model. The function  $\Omega(\mu, \tau)$  has the form in this limit:

$$\Omega(\mu, \tau) \rightarrow 3\Omega_0, \quad \text{where} \quad \Omega_0 = \frac{8\pi G\rho_0}{3\tilde{H}^2}.$$

The function  $\Omega(\mu, \tau) = \text{const}$  in the limit (32) and it is in accordance with condition  $f^2(\mu) = 1$  for the using solution.

## 6 Resultes and Conclusions

This article is dedicated to the development of the Bonnor's solution (9) on the ground of the results has been obtained in [3]. The simplest nonhomogeneous LTB model (13) - (14) generated by the initial conditions (8) and (10) is studied and the follow results are obtained:

- The time of collapse (15) defined by the initial condition  $\dot{\omega}_0(\mu)$ ;
- The expression of the initial conditions through density and Hubble function given at time  $\tau = 0$  are persented in (26) and (27);
- The example of the initial condition with density not dependent on cordinate  $\mu$ , but sutisfyed the FRW equation (23) is shown. The criteria of homogeneouse of the matter distribution in the present solution is shown in (31);
- It is shown that the LTB solution (13) - (14) strives for FRW for time from (32);
- The parameter  $\nu(\mu)$  from (27) defined by the initial conditions  $\omega_0(\mu)$  and  $\dot{\omega}_0(\mu)$  shows the difference between LTB and FRW models;
- The particulary case of initial conditions  $\dot{\omega}_0(\mu) = \text{const}$ ,  $\nu(\mu) = 0$  generat the FRW model;
- The time of collapse in the FRW model is not dependent on  $\mu$ , so all matter fall down to the centre simultaneously.

## 7 Acknowledgements

I'm grateful to Prof.Arthur D.Chernin and Dr.Yuriy Barishev for encouragement and discussion. This paper was financially supported by "COSMION" Ltd., Moscow.

## References

- [1] R.C.Tolman, Proc.Nat.Acad.Sci (Wash), **20**, (1934).
- [2] W.B.Bonnor, MNRAS **159**, 261 - 268 (1972)
- [3] A.Gromov, gr-qc/9606068
- [4] Landau, Lifshits, "The Field Theory", Moscow, "Nauka", (1973).
- [5] Gurevich and Chernin, "The Introduction into Cosmology", Moscow, "Mir", (1978).
- [6] H.B. Dwight, "Tables of Integrals", NY, (1961).
- [7] G.A.Corn, T.M.Corn, "Mathematical Handbook", McGraw-Hill Company, NY, (1968).